

# PHYTOPLANKTON OF LAKES IN THE MUSKOKA-HALIBURTON AREA

L. Nakamoto, L. Heintsch, and K. Nicholls

DATA REPORT DR 83/8

QH 545 .A17 P49 1983 copy 2 MOE

# DATA REPORT SERIES

The data presented in this report were collected by staff of the Water Resources Branch of the Ontario Ministry of the Environment as part of the Lakeshore Capacity Study or the Acid Precipitation in Ontario Study. This unreviewed report does not necessarily reflect the views or opinions of the Ontario Ministry of the Environment

QH 545 .A17 P49

Phytoplankton of lakes in the Muskoka-Haliburton area / Nakamoto, L. 79869

# PHYTOPLANKTON OF LAKES IN THE MUSKOKA-HALIBURTON AREA

L. Nakamoto, L. Heintsch, K. Nicholls

Plankton Taxonomy Unit,

Limnology Section,


Water Resources Branch,

P.O. Box 213,

Rexdale, Ontario, M9W 5L1

- : Ont. Min. of Environment

DATA REPORT DR 83/8



#### PREFACE

The unpublished Data Report Series is intended as a readily available source of basic data collected for lakes and watersheds in the Muskoka-Haliburton area of Ontario. These data were collected as part of the Lakeshore Capacity Study and/or the Acid Precipitation in Ontario Study.

The limnological portion of the Lakeshore Capacity Study (1975-81) was initiated to investigate the relationships between lakeshore development and lake trophic status in low ionic strength Precambrian lakes. The Acid Precipitation in Ontario Study (1979-present) was initiated, in part, to investigate the effects of the deposition of strong acids on aquatic and terrestrial ecosystems in Ontario. The primary findings of these studies have been and will continue to be published as reviewed papers and technical reports.

#### **ABSTRACT**

Phytoplankton biomass of the dominant classes and genera is presented for 16 lakes in the Muskoka-Haliburton area of south-central Ontario. Biomass (or biovolumes) is given both as an annual maximum and as seasonal plots for the May to November periods of 1976 to 1979.

Nakamoto, L., L. Heintsch, and K. Nicholls. 1983. Phytoplankton of lakes in the Muskoka-Haliburton area. Ont. Min. Envir. Data Report DR 83/8.

# Introduction

An objective of the phytoplankton component of the trophic status section of the Lakeshore Capacity Study was to determine if lakewater phosphorus concentrations and other physical-chemical variables were related to characteristics of the lakes' phytoplankton. Of special interest were total phytoplankton biomass (cell volume) and taxa of particular relevance to efficient food chain function (e.g. small unicellular forms) and others which are known to be very inefficient in aquatic food webs and which also impair aesthetic enjoyment and water use (e.g. large, colonial and toxic blue-green algae).

The rationale, objectives and relationship of this investigation to other components of the Lakeshore Capacity Study and to other published studies is presented in more detail in a report nearing completion. The purpose of this Data Report is to provide the most important part of the phytoplankton data set from the four-year (1976-79) study of the 16 Muskoka-Haliburton lakes in a form (tables and graphs of dominant genera and classes) which may be utilized by others undertaking limnological investigations of these lakes or other lakes in the area. The locations and morphometry of the 16 lakes for which phytoplankton data are provided were given in Nicolls et al. (1983). For comparative purposes, data from a eutrophic hardwater system (Bay of Quinte near Belleville) and an acidified Sudbury area lake (Clearwater Lake) have been included in part of the data summary.

### Methods

Phytoplankton samples were collected as volume-weighted composites of the euphotic zones. Full details were given in Scheider et al. (1983).

Samples were fixed in the field with Lugol's iodine solution (containing glacial acetic acid) and returned to the Toronto Laboratory for preparation and analyses. Details of sample analyses (using inverted microscopes) were given in Nicholls and Carney (1979). Subsamples (concentrated) of all collections have been catalogued and retained (preserved with formalin) in the Taxonomy Unit's permanent repository.

The next 24 pages (pp.3-27) contain a listing of dominant genera arranged by class. The numbers indicate the arithmetic mean of the three highest biomasses (or biovolumes) for the year in  $\text{mm}^3/\text{m}^3$ .

Note: Chrysochromulina spp (Prymnesiophyceae) have been included in the Chrysophyceae for convenience and not for any taxonomic reasons.

Class	Page
Cyanophyceae/Dinophyceae	3
Cryptophyceae/Euglenophyceae/Chrysophyceae	4
Chlorophyceae	5
Bacillariophyceae/Xanthophyceae	8
Cyanophyceae/Dinophyceae	9
Cryptophyceae/Euglenophyceae/Chrysophyceae	10
Chlorophyceae	11
Bacillariophyceae/Xanothophyceae	14
Cyanophyceae/Dinophyceae	15
Cryptophyceae/Euglenophyceae/Chrysophyceae	16
Chlorophyceae	17
Bacillariophyceae/Xanthophyceae	20
Cyanophyceae/Dinophyceae	21
Cryptophyceae/Euglenophyceae/Chrysophyceae	22
Chlorophyceae	23
Bacillariophyceae/Xanthophyceae	26
	Cyanophyceae/Dinophyceae Cryptophyceae/Euglenophyceae/Chrysophyceae Chlorophyceae Bacillariophyceae/Xanthophyceae Cyanophyceae/Dinophyceae Cryptophyceae/Euglenophyceae/Chrysophyceae Chlorophyceae Bacillariophyceae/Xanothophyceae Cyanophyceae/Dinophyceae Cryptophyceae/Euglenophyceae/Chrysophyceae Chlorophyceae Bacillariophyceae/Euglenophyceae/Chrysophyceae Chlorophyceae Cyanophyceae/Dinophyceae Cyanophyceae/Dinophyceae Cyanophyceae/Dinophyceae Chlorophyceae/Euglenophyceae/Chrysophyceae Chlorophyceae/Euglenophyceae/Chrysophyceae Chlorophyceae

1976	[Clearwater]	Basshaunt	bniwgia	Blue Chalk	Виск	Сиир	Crosson Dickie	นอุเอ	reather	Harp	Jerry	Little Clear	Red Chalk (east) Red Chalk (main)	Solitaire	Мајкег	[Bay of Quinte (B)	
CYANOPHYCEAE						Ø	max	. (mm <sup>3</sup> /m <sup>3</sup> )	/m <sub>3</sub> )					/ 5			
Anabaena Aphani zomenon		ω	4 п	22	и	He		П	2.0	42 5 0		0.2	22	14		5826 2679	
or Gomphosphaeria	2.5	0.5	1154		30 30	2 0 ° 0	6000	12 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		16	227	0 1 m u	33			30 118 290	
Dactylococcopsis ∝ or Rhabdoderma Gloeocapsa Gloeothece Lyngbya		υ • 0	0.5		H	7 0	0			35	35	23	7 2	0	5	393	
Marsoniella Merismopedia Microcystis Oscillatoria		0 .0	B -	12.07		2	3 179		0,3	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	Н 4	0.5	0.0	000	0.3	48 591 2430	-3-
Phormidium Pseudoanabaena Radiocystis Raphidiopsis							1	m north		142.2						322	
Synechococcus Synechocystis Unid. Cyanophyceae	0.5	Н					4	-					7				
DINOPHYCEAE Ceratium	Ecre							ele aus				0.2	106				
Glenodinium Gymnodinium Peridiniopsis	21	0.5		<b>C</b> 3	m	95	8 147	7 43	19	26	17		m		2 0.3	. 6	
Peridinium Unid. Dinophyceae	893	44		4	36	62 1	124 78	ω <sub>0</sub>	298	32	m		ω		8	383	

	A
_	/1

			-4-	-	
[Bay of Quinte (B)		1112 43 83	252	22	36
Мајкег		12		e .0	0.3
Solitaire		56		mm ⊢	7
Red Chalk (main)		104		1 65	13 0.2
Red Chalk (east)					and trigger (mental)
Little Clear		128		NW	21 21
Jenry		166		Н 0	0 0
Нагр		132		HH 24	0.2
Gullfeather	(mm 3/m 3)	179	m	ИW	39 3
nələ	( mm )	15		7	0.0
Dickie	max.	112	59	0 00 w ruru	94
nosson	I	117		0.2	4 1
Сһир		131 28 1	18	H4, 00	e . 42
Buck		7 H4	9	H4	45
Blue Chalk		61 83 0.5		ww.	0.5
bniwgia		21 0 .5 0 .5	m	O • O	
Basshaunt		123		Н	
[Clearwater]		00 00 00 00 00 00		· 0	0 0
976	RYPTOPHYCEAE	Chroomonas Cryptaulax Cryptomonas Katablepharis Rhodomonas Unid. Cryptophyceae	UGLENOPHYCEAE Euglena Lepocinclis Phacus Trachelomonas Unid. Euglenophyceae	HRYSOPHYCEAE  Bicoeca Bitrichia Chromulina (Monochrysis) Chrysoamoeba Chrysidiastrum Chrysochromulina breviturrita	

		1				-5-				
(8)	Eay of Quinte		153	1 45	17	110	9 3 3		15	27
	Мајкег		101	0.3		16 51 0.3	~	45	-1	
	Solitaire		т	0.3		012				
(u	Red Chalk (mai		19	63	9	49	15	73	12	
(7	Red Chalk (eas									
	Little Clear		60)	1 68		14 56 11			11	
	ენის		v	0.3	2	16 1 1	П 4. С		10	1
	Harp		122	1 224	0. 4. 1.	5 72 1 27	9 n	399	0	4
	reather	(mm <sub>3</sub> /m <sub>3</sub> )	30	0.2	17	82 103 4 113		31	10	
	gjeu		9	5 0 3	. 2 H	4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4		0	11	
	Dickie	max.	416	0.5	0	584 3443 0.2		80		9
	Crosson		0.2	. 17	1 0.2	0 99	2	01	89	
	dudo	-	308	3 1 5 181	11	3 152 2 0.2 2 5.6 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	7	82	0 12	2
	Buck		174	0.3		2 2 2 3 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	51 24	15	17 10	0.2
	Blue Chalk		47	0.5	m	0 0	ιΩ			10
	bniwgia		21	0.5	0.5	5 5 5 9 66 5 1 4 5 1 4 4		12	14	0.5
	Basshaunt		2 0.5	വയ	ıΩ	0.1			11 5	. 10
	[Clearwater]		. 72	16	0.5	0 184 0 7			П	0.5
	1976	CHRYSOPHYCEAE (Cont'd)	Desmarella Dinobryon Epipyxis	Kephyrion Mallomonas	Pseudokephyrion Rhizochrysis Salpingoeca Spiniferomonas	Synura Synura Uroglena Unid. Chrysomonads Unid. Chrysophyceae Chrysophyceae	CHLOROPHYCEAE Actinastrum Ankistrodesmus Arthrodesmus Asterococcus	Binuclearia Botryococcus Carteria	Characium Chlamydomonas	Chodatella Closteriopsis Closterium

							-6-				
[Bay of Quinte (B)		268	9	54	-	œ	163	<u>(</u> 0.	26 448	162	358
Malker		4 m		.2	0.2		4		e 0	ю .	0.3
Solitaire				.2 0	1 0		26	0.2	4	26	
Red Chalk (main)		თ ⊢	rI	7 0	н		61	J	<del>г</del>	11	
Red Chalk (east)											
Little Clear		3	7		т		7	2	0	m	
ીલ્પાપ્		Н.	0	н			m	0	-	14	4
qrsH		16	m	4			36	m .	m m .•	7 1	00
mentsel[[u2	/m <sub>3</sub> )	52	24	4	п		45	•	0	49	
นอเา	( mm )	119		r1	2		48	0	<b>4</b> <sub>1</sub> .	0.5	
Dickie	max.	1 1 23	ω	· 	1 55	•	16	0.2	m v	78	
Crosson	m	<b>4</b> ε		m	्रात		4	0 1	16		
Chub		e E	2	m	e		24	нн	1 0.	15	, <del>erri</del>
риск		7		н	.2		4		, <b>–</b>	10	
Blue Chalk		10.0		2	0		ю	⊢	н	22	4.
bniwgia Wiod2 owid		9 4			rel		3 23		-22	13 2	
		-1 -1 -2		5 0.5		m			0.5		
Basshaunt		A: <del>1</del>	7	5 0 5		,	Ŋ			5 0 5 5	
[Tearwater]		2 4		0 . 5			0.5			345	
1976	CHLOROPHYCEAE (Cont'd)	Coccomyxa Coelastrum	Crucigenia Cylindrocystis	Uicellula Dictyosphaerium Dimorphococcus	Echinosphaerella Elakatothrix Euastrum	Eudorina Franceia Geminella	Gloeocystis Golenkinia Gonium	Gyromitus Kirchneriella Koliella Lobomonas Micrasterias	Monomastıx Monoraphidium Mougeotia Nephrochlamys Nephrocytium	Oocystis Pandorina Paramastix	Paulschultzia Pediastrum

	1							_	7-								
(B) SaniuQ To vs8]				m	9	152	7		H		78		30	7 1		-	57
Malker			m		m	-							c				
Solitaire					Ó	.2	Н		23								0
Red Chalk (main)			2		166	1 0			0								16
Red Chalk (east)					-												
Little Clear		Н							0.5				0.2				32
ენĸĸλ					Н	0.2			7								O
Harp					က	0 9			4		10		ر د د	e. 0			43
redtael[[u2	m³)				2	9	-		20				2				21
นอโอ	( mm 3 /m 3)				4	0.3							41				41
Dickie	max. (	П			·0	5			-1								10
Crosson	me				9				0.5					-			m
супр					4	щ			0.5	7	27		7				9
Виск							H				42	:					Ŋ
Blue Chalk			-		9	Н	Н		11	4	13	35					13
bniwgia					0.5		0.5										2
Basshaunt					0		0.50										0
[Clearwater]						0.5	O				152	)	7			4	4
						J								*.			
																	a
	nt'd)			ja B						v	,						Chlorophyceae
	(CO	nas	ema	phaer	as	cus	lamys ria	ldia	mystis	OSMA Zonsi	Sium	Smms	ccus	m a	e B		lorop
	HYCEA	Pedinomonas	Phacotus Planctonema	Planktosphaeria Polytoma	Pteromonas Quadrigula	Radiococcus Scenedesmus	Schizochlamys Schroederia	Scourfieldia Selenastrum	Sorastrum Sphaerocystis	Sphaerozosma Spermatozonsis	Spondylosium Ctauractrum	Staurodesmus	Stichococcus Tetraëdron	Tetrastrum Treubaria	Trochiscia	Westella Xanthidium	Unid. Ch
1976	CHLOROPHYCEAE (Cont'd)	Ped	Pha Pla	Pla	Pte Qua	Rad	Sch	Sco Sel	Sor	Sph	Spo	Sta	Sti Tet	Tet	Tro	Wes	Uni
. 51	. I 5																

[Bay of Quinte (B)			609		254	625	450	431424	1 23	352	6027	73
мајкеr			35		108			m	0.3	11	ŭ	1 8 1 2
Solitaire			12		26		Н		0.2			mi
Red Chalk (main)			134		1939	-	37		0.3	0 3	0.3	81 0.3
Red Chalk (east)												
has Clear		0.3			171				1	Н	23	149
ენსა					72	2	4			10		on '
Harp			48		107		45	0	က	750	56	29
reathealling	( mm <sub>3</sub> /m <sub>3</sub> )		319		20	7		306		90	Ŋ	m
บุลโอ					137			•	01.01	_1	1	m m
Dickie	max.		97		23			223	0.2	111	21	0.3
Crosson	-				C)	90		Н		16		1 266
СһиЪ			24		28		٦	7	7	184	4	164 865 4
Bnck			16		155				2		М	0.2
Вјие Срајк !			20	H	217		Н		Н			253 4
bnîwgi8 j		٠. ت	4				0.5	29		177	1	103
Basshaunt												Φ
[Tearwater]		32	0.5	H	7	. ~			H		Н	0.0
76	CILLARIOPHYCEAE	Achnanthes Amphora	Asterionella Atthova	Cocconeis Coscinodiscus	Cyclotella Cymbella	Diatoma Eunotia	Fragilaria Frustulia	Gomphonema Melosira Meridion	Navicula Nitzshia	Rhoicosphaenia Rhizosolenia Skeletonema	Stenopterobia Stephanodiscus Surirella	Synedra Tabellaria Unid. Diatoms NTHOPHYCEAE

9

Isthmochloron Ophiocytium Pseudopolyedriopsis

											-9-								
[Bay of Quinte (B)		8284	1	33	154			94	23	119	647		16			0.2	861	180	207
мујкек		11		<b>~</b> ⊢				7		0.3	•			c				11	23
Solitaire		9		13	4 0	υ Ω					m							19	· 8
Red Chalk (main)		25	3 0.	76 24	16	0		9	4	ហ	7		0.2	r	-1	12	1	10	40
Red Chalk (east)			28	33		7		m	12	12	10		0.3 0	r	า	0	1	30	135
raefo efttid		20		31	ω	ر ک		2	ω	11	4 <sub>1</sub> 0		H	,	7 <del>4</del> 0		7	1	56
Jerry		48	4	95	39	94	7	17	ω	<b>-</b>	7			,				6	24
Натр		41	.55	55 14	25	20	21	00	4	35	m							46	83
redifeail[lu2	/m³)		2 1	4°W	0.3				H	7	0.2			•	-1				15
นอโอ	(mm <sub>3</sub>		•	00.5		,		0.3		0.3			· · · · · · · · · · · · · · · · · · ·		0.2				
Dickie	max.		.2		30	20	٠		87	H			0,3					0	174
Crosson	4		1 0	16	28	58	-1		30	0.2					Н			7	18
Сћир		20	H	<b>ω</b> 4	12	127		0.2	10	-	0.2		0.2	•				46	0.3
Bnck		2	e,	62	2			٦ ا	m		-		·· . · ·					m,	172
Blue Chalk		27	0	13	19	m		7	7	Н				0.2			0.2	5	14
bniwgia		7	r-j			-			2	9	0							Н	m,
Basshaunt			Ŋ		.2			0.3	5						0°3				
[Clearwater]				H	1 0			0			18		•		7			106	164 221
,	,				eria	ma.													
					or Gomphosphaeria	or Rhabdoderma													
					Gomphe	r Rhal													
						8									eae				9
		2	= m	αυ <i>ν</i>	rium	copsi			ص بــ ص	. s	ia	aena	S I	tis	Unid. Cyanophyceae	ε	S	EE	nium Dinophyceae
	CEAE	ena	caps	othec	sphae	lococ	thece	/a :	noped	cysti	lator	oanab	diops hococ	hocys	Cyan	<b>EAE</b> diniu	ium psali	diniu diniu	m open
7.7	CYANOPHYCEAE	Anabaena	Aphanocapsa	Aphanothece Chroococcus	Coelosphaerium ∝	Dactylococcopsis Gloeocapsa	Gloeothece	Lyngbya	Marsoniella Merismopedia	Microcystis	Oscillatoria Phormidium	Pseudoanabaena	Raphidiopsis	Synechocystis	Unid.	DINOPHYCEAE Amphidinium	Ceratium Diplopsalis	Glenodinium Gymnodinium	Perid Unid.
1977	CYA															DIN			

	1 1		-10-			
[Bay of Quinte (B)		5 3 1244 20 70	225 0.3 1 55	87	Н	2
Ма]кег		41 6 11 0.3		0 0	н.	12
Solitaire		67 6 38 0.2 (	н	m	7	
Red Chalk (main)		62 10 7 0.2 (	70	<b>ч</b> 4 ч	0.5	rv
Red Chalk (east)		0.2 53 11 15	10 6 10	H & H	4 5 6	192
Little Clear		0.3 118 6 68	Н 4	U 9	D. 3	
ენსა		405 18 32	· o	2 1 2 5	1 6	
Harp		207 4 9 18		11	9 7	36
red lifeather	/m 3)	244 2 10 5	42	н е н	о 1	Н
ຕອເລ	(mm <sub>3</sub> /m <sub>3</sub> )	260 1		. ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	7	
Dickie	max.	82 14		m 0	7 7	73
Crosson	5	2 4 2 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2 4 2		H	0 3	н
Сһиь		234 11 7 0.3	32	0.24	7	72
Buck		66 7 16 0 5 6	10	0.2	4 .0	2 0
Blue Chalk		0.2 122 40		757	<b>ω</b> κ	w.
bniweid		8 7 7		7 7	7	н
Basshaunt		57		. 0	0.5	
[rearwater]		187 1 0.3				
1977	RYPTOPHYCEAE	Chroomonas Cryptaulax Cryptomonas Katablepharis Rhodomonas Unid. Cryptophyceae	Euglena Lepocinclis Phacus Trachelomonas Unid. Euglenophyceae	(Monoc ba trum mulina	Chrysochromulina spp. Chrysococcus Chrysolycos Chrysosphaerella coronacircumspina	

			-11-		
[(8) Sturing to Vall	0.38	16	34	11	130 205 205 237 237 3
Malker	101	20	110 115 116 116		_
Solitaire	87 H	0 2	120 1 120 1 77	0,3	7 9
Red Chalk (main)	22 1 0		21 21 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	-	31 0 . 5
Red Chalk (east)	69 7 5 11 7 5 11	γ) •	524 524 33 33 33		
Little Clear	26	0 1	12 127 2 0.3		10
декку	130	7 7 7 7 8	68 109 1 1 C	12	2 04 6
Harp	ω m α	20 21 12	140 57 1 3 66	31	13 8
79d1feal[[u2] € ∃	23 21	5	34 16 3	23 3	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
mm % m Glen Gullfeather	0,0	7	41 - 8		0 . 2
B Dickie		104 1 6 0.3	1200 18 94 0		164
nosson	2 L L	3	33 1 33 2 33 3	0 6 4	8 8 2
СРИР	101 2 0	59 1 2 0 0 2	509 29 33 39	16	12
Виск		28 1 0.3 (	24 1 51 0.3	0 . 5	Φ
Blue Chalk		4 4 4	138 444 90 2 23	12	6 6
bniwgia	173 0.3 0	7 7	18 1 43 1 40	39	12 3
Basshaunt		4 .0	28 15 14		e e
[Clearwater]	35	9	232 232 7		40
ont'd)	•	uo s	Stelexomonas Synura Uroglena Unid. Chrysomonads Unid. Chrysophyceae Chrysophyceae cysts	S	
AE (C	, פר ה ה	as as phyri ysis eca eca	rysom rysop yceae	AE rum desmu smus	ria ccus m monas a la opsis
PHYCE	Desmarella Dinobryon Epipyxis Erkenia Kephyrion	Mailomonas Ochromonas Pseudokephyrion Rhizochrysis Salpingoeca Spiniferomonas	Stelexomonas Synura Uroglena Unid. Chryso Unid. Chryso	OROPHYCEAE Actinastrum Ankistrodesmus Arthrodesmus	Binuclearia Botryococcus Carteria Characium Chlamydomonas Chlorella Chodatella Closteriopsis
1977 CHRYSOPHYCEAE (Cont'd)	Des Din Epi	Mail Och Pse Sal Spi	Synura Synura Uroglo Unid. Chryse	CHLOROPHYCEAE Actinastru Ankistrode Arthrodesm Asterococc	Bir Bot Char Char Char Char Char Char Char Char

1977

	1 1							_	13-						
[Bay of Quinte (B)		15	4	m	2	172	15				222	28	22	39	18
мајкег			0.3		1	Н									7
Solitaire						0.3	0.2								12.
Red Chalk (main)		.3	0.3		'nΩ	3	0.2 (	H	2		11	H			7
Red Chalk (east)		0.2 0	7 0		00	9	0,3 (		2		7	∞ -	4	4	31
Little Clear		O		0.2	7	c	0.3 (					<del></del> -			்ம
Yarel		Н	7	J	S	(L)		0.2		0	0.2	H 0	• •		러
quaH		r	-1		<u>ه</u> .	ന്നു.	0.2	0	03	m	7	·		ന	NW
reather	(mm <sub>3</sub> /m <sub>3</sub> )		7	0.3	0.3	9	0.2		2		0.2	<u>ر</u> د	0		26
นอโอ	(mm					0.2	Н								0.3
Dickie	max.				Ŋ	13 0			7		18	ru c	0		4
nosson					က	Н						7	0.5		H '
СһиЬ				0.3	9	m	0.3			0.3	100	7	0.3	٠	Н
Виск				J	0.3	36	0.2			0.3		(	7.0		m
Blue Chalk		0.2	က		5	Н	Н				65	m (		-	4
bniwgia					က	Н			Н		2	H (	7.0		7
Basshaunt					0.2		0.3		0.3						H
[rearwater]						0.3		,	0		14				Н
1977	CHLOROPHYCEAE (Cont'd)	Pedinomonas	Phacotus Planctonema	Planktosphaeria Polytoma	Pteromonas Quadrigula	S S	Schizochlamys	Scourfieldia Selenastrum	Sorastrum Sphaerocystis	Sperma tozopsis Spondylosium	Staurastrum Staurodesmus	Stichococcus Tetraëdron	letrastrum Treubaria	Trochiscia Ulothrix	westella Xanthidium Unid. Chlorophyceae

	1	l								-1	4 –								
[Bay of Quinte (B)		н	714		30 140	1333	1 6	324	26024		43		59 26	5132	1350				
Malker			35		41	•			•				11		18				1
Solitaire			31	,	201									13	0.3				
Red Chalk (main)		2	52		1534		m r	<b>-</b>	• *	m	സ്ധ	۱. د	0		7 (				
Red Chalk (east)		0.2	46		1160		0 0	•	37		0 HH		21	'n	6 139 1				
rabil bittid		0.30	12		477 1		c	<b>.</b>					9		7) L				
Jerry		0	17		570 4	m	: ;	ý	. •	. : 1			1117	256	134			Н	
Нагр		0.3	32		422		·	) ,		,	-		841 1	15	41 :				
reather	/m <sub>3</sub> )		148		47	:			299	. (	nΗ		116 8	14	17				
นอเอ	(mm <sup>3</sup> /m <sup>3</sup> )				81		÷ , .								7				
Dickie	max.		82		m	,	-	4	158	4.			_	7	3				
Crosson					, m		7			,		25		0.3	81 ]				
Сһиь			30		16		y	)	89		0.3		97	5	8				
Виск			4		30										il c				
Bine Chaik			26		160	Ŋ.	0	•		Ų	0H '	ı	-		4				
bniwgia		0.3			30		•	4	65	c		. 1	17	17	892				
Basshaunt		O	ω		0°3						,				77				
[Tearwater]					m					, (n	n ·			7	0.3			17	
						-													
																			is
977	ACILLARIOPHYCEAE	Achnanthes Amphora	Asterionella Attheva	Cocconeis	Cyclotella Cycholla	Diatoma	Eunotía Fragilaria	Frustulia	Melosira	Meridion Navicula	Nitzshia Pinnularia	Rhoicosphaenia	Skeletonema Stenonterobia	Stephanodiscus Surirella	Synedra Tabellaria	Unid. Diatoms	NTHOPHYCEAE	Isthmochloron Ophiocytium	Pseudopolyedriopsis
	ACILLARIOPHYCEAE	Achnanthes Amphora	Asterionella	Cocconeis	655	Diatoma	Eunotia Fragilaria	Frustulia	Melosira			Rhoicosphaenia	Skeletonema Stenontemokia	iscus	0	Unid. Diatoms	ANTHOPHYCEAE		Opinocytium

- 4		
_	ו ו	-

						-3	L5-				
[(8) of Quinte (8)]		2287	84- 804-	121	86	3 5 9 9	257		0.5	252	54 236 369
Malker		Н	0.42	0.00	Н	ო			0.2		51
Solitaire		0.2	21,	32		16	Н	Н			20
Red Chalk (main)		44	79 67 72	57	m	20	0.2		21	42	7.5
Red Chalk (east)		25	16 89 69	23	ហ	22	0.3				
Little Clear			0.3	14	0.2	40	1 5				111
วิธนะวิ		9 7	7	215		88					37
Harp		180	37 53 19		52	69	-				175
Gullfeather	/m <sub>3</sub> )	0.2	0.244	7	00.3	0.2	•				213
นอเอ	( mm <sub>3</sub> /	6	23 25	4		-	54		-		259
Dickie	max		.0 .0	ო ⊢	-	ထဖ					304
Crosson			H 62	2 H		19					24
Сћир	-	4	0.2	m 7		0.0	0.2				155
Buck		0.2	0 54 m	H			0.3				176
Blue Chalk		23	36 15 29	24 1	œ	44	7 7			0 5	41
bniwgia		m	040	0.55		0.2	7			Ü	12
Basshaunt		нε	23		0°3	16			0.2		30
[Clearwater]			00.0			(	.0		0.5		211
	-			or Gomphosphaeria ∝ or Rhabdoderma						,	
1978	CYANOPHYCEAE	Anabaena Aphanizomenon		rium ∝ copsis	Gloeocapsa Gloeothece Lyngbya Marsoniella	Merismopedia Microcystis	Phormidium Pseudoanabaena	Radiocystis Raphidiopsis Synechococcus Synechocystis	Unid. Cyanophyceae	Ceratium Diplopsalis Glenodinium	Gymnodinium Peridiniopsis Peridinium Unid. Dinophyceae

			1				17	-							
](	Bay of Quinte (B		111	2 70			9		15		ហ	53	90	4	33
	Malker		41	1 27	-	H 2.	2	16	36		10	ſΩ	Sin		
	Solitaire		4	.5.0	•	1.50	2 2		2.5		10	27	0 m		
	Red Chalk (main)		111	33 0	1 0	1 0 1 0		42	0 8	)	4	12	7	e . 0	0.3
	Red Chalk (east)		230 ]	63	ក់ (		7	92	1 23		0 0 0	43	6	0	Ü
	TESED SETTIA		37 2	1 12	2	0.2		241 	49		10	6 0	9		
	ენას		139	3		1 1		96 61	33 33		3		œ	Н	6
	Нагр		15	4	r-l 1		2	87	53		e 0 e	49	40		13
	Gullfeather	/m <sub>3</sub> )	52	30	•	0	30		38		m	146	22		
	uə[g	( mm <sub>3</sub> /	7	83		7 77		93 1	1 26 1		0	14 1	4		
	Dickie	max.	E = 1	28 0 39		2 00 C	60	L <sub>2</sub>	29		1 68	96	11		
	nosson	m	26 66	17	7. 0	0	8	11 57 11	m 7		. w		 14		
	Сһиь		П 4		0	വ നന	m	2	00		41	59	37 0.		
	Bnck		2 42	4	- L	0	Н	ORI					2		
	Blue Chalk		5 12	2 1 7 32		3 0,3			1 27 22		1 15 0.3	2 114	10		
	bniwgia		N	Н		1 0.3		17 30 57 300	15 2			7 0.2	2 1		
	gasshaunt		0	1 5 4	m L	0	-				2 0.2	43	6		
	[Tetewrse[]]		47 5	,	° °				.2 0.2 17 9		0.2		41		
	[motermeo[J]		4,				-	0 4 28	0.2			0.2	4		
		(p,						ds	eae						
		(Cont			rion s	nas s		omona	chrysophyceae phyceae cysts		mus s	v	as	•	s
		CEAE	ella yon is	ion	kephy hrysi	eromo omona		Chrys	chrys	CEAE	rodes desmu coccu	coccu ia	domon 113	e] a	riops
	m	CHRYSOPHYCEAE (Cont'd)	Desmarella Dinobryon Epipyxis Erkenia	Kephyrion Mallomonas	Pseudokephyrion Rhizochrysis	Spiniferomonas Stelexomonas	Synura	Unid. Chrysomonads	Unid. Chrysophyceae Chrysophyceae cysts	CHLOROPHYCEAE	Actinastrum Ankistrodesmus Arthrodesmus Asterococcus	Botryococcus Carteria Characium	Chlamydomonas	Chodatella	closteriopsis Closterium
	1978	CHRY	0000	ΧΣĊ	<u>a</u> æ <i>v</i>	SON	S)	) <b>)</b> :	50	CHLO	4444	a a C C	っここ	១០០	ວີ

				0.00	-18	27	2 3 7		23	9	ÖM I	_ ,	ത
[Bay of Quinte (B)		130	8 221 0.2	0.2		HN	• • • • • • • • • • • • • • • • • • • •		7		9.0		22
Malker		13	H	0.3		16	m				2		
Solitaire		0 333.1		7		43			0.5		0		
Red Chalk (main)		8 118 47		-	,	316	1				10		
Red Chalk (east)		7 29	Ŋ	ю		158	7			2 0	18		0.2
Little Clear		31 1	-1	00.5		51	0.22		0.3		71/2		0.2
<b>ე</b> 6ಒւλ		111 22 4	4	0.3 6.4		32	4 .0		m		14		
Harp		33	6.3	0.2		38	7 0.3	Ŋ	2	7	26	7	4
redtael[[u2	/m <sub>3</sub> )	10 51 13	0.2	0.2		50	0.5		13	2	35		0.5
uəլg	(mm <sub>3</sub>	09	N.			147	0.2		0.5		13		
Dickie	max.	21.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7	11 0	0		19	0.2		0.5		14	7	
Crosson	=	0	ო	Н		22					က		
dud		2 22 13	15	0.2		71	26 2 0.5		0.2	ശ	147	*	0.2
Виск		0.2 4 15		.2		2			0.2		ω.		
Blue Chalk		60 53	m	0.50		64	0.2			0.3	14		
bniwgia		6 28 0.3		0.3 0		92	Ŋ				18		
Basshaunt		0.2 23 C	e 0			9	0.2		0.3		m	0.3	
[Teatewres[]]			7 E			4	51		ပ		176	U	
		·									• •		
1978	CHLOROPHYCEAE (Cont'd)	Coccomyxa Coelastrum Cosmarium Crucigenia	Oyling Ocysels Dicellula Dictyosphaerium Dimorphococcus	Elakatothrix Euastrum	Eudorina Franceia	Gloeocystis Golenkinia	Gyromitus Kirchneriella Koliella	Lobomonas Micrasterias Monomastix	Monoraphidium Mougeotia	Nephrocytium Octobrium	Oecystis Oocystis Pandorina	Paramastix Paulechultaia	Pediastrum

[Bay of Quinte (B)]		27		14	213	Ŋ	-3	19-	Н		22	m		67
Malker					П						-!	0.2		C)
Solitaire						m •			0.2			0		ო
Red Chalk (main)				3	-	0			0.2	33	-			
Red Chalk (east)					2	.2			0	7 7	m -1			9
rittle Clear				7	4	1 0,			C	0 9	2 10	0		
า กับ เกิด เกิด เกิด เกิด เกิด เกิด เกิด เกิด				4	9			<b>-</b>	.5		100		m	Н
Harp				7	12			(	0		2.0		0.5	П
rediael[lu2	m 3 )	Ŋ		0.5	25					0 / 7	0 3 0		O	2
gjeu	( mm <sub>3</sub> /m <sub>3</sub> )			5	m	2	9.				m			14
					10 0.		4		2		w 010	2 .		m
Dickie	max.			2 14	5 1				0.2	137	Ó	0		0.2
Crosson				c	3 0.	2			m	4	7			L 12
Chub	-				2	.2 0.			0	.2 6		0.2		0.30
Bnck				0.5		0				0	<del></del> !	0	Н	3 0
Blue Chalk				m	7	0.5				5 0.2		2		
bniwpia			•	9	Φ					9	2	2 0.2		0 3
Basshaunt				0.5	0.2	0.3				6	0.2	0.2		15
[Clearwater]				0.3	1									-1
	·													
1978	CHLOROPHYCEAE (Cont'd)	Pedinomonas Phacotus Planctonema	Planktosphaeria Polytoma	Pteromonas Quadrigula	Kadlococcus Scenedesmus	Schrzochlamys Schroederia	Scourfieldia Selenastrum	Sorastrum Sphaerocystis	Sphaerozosma Spermatozopsis	Spondylosium Staurastrum	Staurodesmus Stichococcus Tetraëdron	Tertrastrum Treubaria	Trochiscia Ulothrix	Westella Xanthidium Unid. Chlorophyceae

	1 1							- 2	20-						
(B) Sy of Quinte (B)		Ċ	391	5	56 97		436	8926	1 23		99	2464	8 6 8 8 8 9		0.3
Malker			29		227	0.3			0.2		11	25	0.5		
eritatilo2		0.2	36		130				-				24		
Red Chalk (main)			40		689					0.2			12 53		
Red Chalk (east)		٦	13		102		П		0.5	ı	2		64		
Little Clear			7		88			2	ני	•	0.3		22		
Jerry			4239		162			9		-	784		74		
Нагр		Н	103		103		ഗ	20	0.2		141	Ω	342		4
red teat [ lua	( mm.3 /m3)		3.1		116			86	0	•	48	0	15		က
gjeu	(mm)				980	0.2							0.2		
Dickie	max.		06		5 94	J		273	_	4	494	16	68 51		
Crosson	Si .		736		က			,			09		228		
Сћир		m	592		34			0.2	<del>≓</del>		110	7	325		
Buck			0.3		327								0 .3 4		
Blue Chalk		H	2		389	-							152		
bniwgia			38		29		23	37			Н		10		
Basshaunt			7		326				0.3	•	Н		79		
[Clearwater]		5			0.3	٠ ر	0.2		0.2				0.2		
1978	BACILLARIOPHYCEAE	Achnanthes	Asterionella Attheya	Cocconeis Coscinodiscus	] a	Eunotia	Fragilaria Frustulia	ща	Meridion Navicula Nitzehia	Pinnularia Phoioschaonia	Rhizosolenia Skeletonema	Stenopterobia Stephanodiscus Surirella	Synedra Tabellaria Unid. Diatoms	XANTHOPHYCEAE	Isthmochloron Ophiocytium Pseudopolyedriopsis

	2	7	
-	1	- 8	-

	2	2	
_	_/	_	-

					-22-							
[Bay of Quinte (B)		647 33 61					0.3		28	Н		
Majker		0.2 4.2 7.0 0.2			0.2				ro.	Н	381	
Solitaire		3.14.0			J		100.5		72	.5	6	
Red Chalk (main)		0.5 0 53 7					2 1 1 0	Н		1 0	23	
Red Chalk (east)		0.2 60 7 11 0.5			0,5		H H 8		34	0.5	1042	
Little Clear		53 2 2 2 2 0		7	0.2		0.2		m	0.5	51	
Jerry		189 25 24					<u>н</u> н и	-	13		285	
Нагр		84 12 13		ω			0.5		14	2	66	
redisel[[u2	/m3)	69 13 0.2		пн	<u></u>		нню	7	7	0.3	278	
nele	(mm <sub>3</sub>	219 10 48					0,3		19			
Dickie	max.	1022 19 1			30		H H 8		2438	H	10	
Crosson		27 4 4 1		e • 0	H		0.3		4	0.2	7	
Сһиь		93		H	9		446		4 2		252	
Виск		2 64 8 10			0.3		0.3		18	7	97	H
Blue Chalk		8 7 8 1		Н .			нню		9	0.2	15	
bniweia		е 4 г с					0.5		8		23	
Basshaurt		11 8 0.2		Ŋ	က		ਜਿਜ	m	4 4	Н	37	
[rearwater]												
1979	CRYPTOPHYCEAE	Chroomonas Cryptaulax Cryptomonas Katablepharis Rhodomonas Unid. Cryptophyceae	EUGLENOPHYCEAE	Euglena Lepocinclis Phacus	Irachelomonas Unid. Euglenophyceae	CHRYSOPHYCEAE	Bicoeca Bitrichia Chromulina (Monochrysis)	um 1 ina	Chrysochromulina parva Chrysococcus		Chrysosphaerella longispina Chrysosphaerella spp.	Codonosiga Codonosiga Derepyxis

									-2	23-	-											
(8) etniup to vea]		19	,	16 16			Н				09	7	13					r	14	134	Ŋ	4
Mg]ker		106	7 !	1 28	0.5		4	П	229	69	34	Н	38			0	7	12		0		0.2
Solitaire		12 1		17			Ţ			12	28		10			-	<b>.</b> 1,	26		m		
Red Chalk (main)		50	7 1	7	Н	C	n 0	0°9	0	Ŋ	20	П	20			c	า	73	7.0	7-	-	•
Red Chalk (east)		68	ກ	35		0,3	-i 9	77	46	55	104	က	23			r	ń	61		9		
Little Clear		4		<b>н</b> б		_	4	00		2	69	}	31			,	71	15		-		
ปิยหาง		20	ဂ	48	4	•	44	24	84		248	Н	22			20		5		16		.0
queH		347	, C	87	7	(	m m	m⊢	24	11	96		39			ת ע	<b>†</b>	78		1.1		5
rediselling	/m <sub>3</sub> )	62		26	2	0.2	m	ന	53	2	47		134			<b>⊢</b> <	t <sup>*</sup>	107		20	•	
นอเอ	( mm <sup>3</sup> /		•	5	0,3		0.5			П		0.3							7.0	17	•	
Dickie	max. (	64	4	10	m	9 1		2-	145	$\vdash$	Н	7			Н	ן ה	-1	142		18	0.2	56
Crosson	me	21 2		37			m	0.3	9	٦		0.2					4	က		2		
Chub		65	<del></del>	25	24		ო	m ~	185	47	71	7	78		0.2	0.3		97	ກ	16		
Виск		122		4 7			2	070	• 9	28	41	<b>н</b>	42			c	•	16		9		4,
Blue Chalk			ហ	າ ແ	0.3		27	700	37	49	79	<del></del> 1	30			0.2	0	35	0.3	13		0.3
bniwgia		56	0.3	1 05	0.5		2			26	16	0.5	18			0.3	0	18		-		0.3
Basshaunt		29		2 7			m	· 产	24	6	45	0.2	38			c	•	14		00		-
[retemrasf2]																						
,																			•			
1979	CHRYSOPHYCEAE (Cont'd)	Desmarella Dinobrvon	Epipyxis Erkenia	Kephyrion	Ochrononas	Pseudokephyrion	Rhizochrysis Salpingoeca	Spiniferomonas	Synura	Uroglena	Unid. Chrysomonads	Unid. Chrysophyceae	Chrysophyceae cysts	CHLOROPHYCEAE	Actinastrum	Ankistrodesmus Arthrodesmus	Asterococcus	Botryococcus Carteria	Characium	Chlamydomonas Chlorella	Chodatella Cloctonioneia	Closterium


					24-					
[Bay of Quinte (B)		69	43	17 4	31	4 4	23	. 2	60	181
Мајкег		4.0		0 0 . 2	8 E	<b>r</b>	٠.	<b>⊣</b> ,		
Solitaire		1 28 41	Н	0°3	95	0.2		0.3	17	
Red Chalk (main)		0.5 35	0.3	0.5	0.2	0.3			7	
Red Chalk (east)		0.3	7		56	0.3	0.5	<b>⊣</b> :	1	
Little Clear		4 2 9	0.2	ы	203	0 • 5	0.2	C	0.2	
ენისე		7 33 33	0.3	0 0 0	0	0.2	, <del>H</del>	ŗ	27	
Harp		1 50 18	2	11	17		7 '	⊣ (	1 1	7
redifeatilua	(mm 3/m 3)	2 7 13	2		30	<u>ი</u> ო	7	7 ,	1	
เลยเอ		12 1 37			77	0 0	2	4	2,	
Dickie	max.	нен	0.5	0.2	11	m ⊢m	16	r	1	2
Crosson		0.2		0.3	118	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0.2	.(	л <del>н</del>	
qnų	-	3 27 31		0.2	123	12	0.3	- ;	00°3	19
Buck		4 w r	0.5	0.2	15	4	<b>-</b>	N 1	n	0.2
Blue Chalk		13	5	0.5	32	7 11	H (	12	0.2	0.3
bniwgia		0.3			13	0.2		(	7	-
Basshaunt		4			က	H H	0.2	(	ת	
[Tearwater]										
1979	CHLOROPHYCEAE (Cont'd)	Coccomyxa Coelastrum Cosmarium Crucigenia Cylindrocystis	Dictyosphaerium Dimorphococcus Echinosphaerella	Elakatothrix Euastrum · Eudorina Franceia	Geminella Gloeocystis Golenkinia Gonium	Gyromitus Kirchneriella Koliella Lobomonas Micrasterias	Monoraphidium Mougeotia Nephrochlamys	Nephrocytium Oedogonium	Pandorina Paramastix Paulschultzia	Pediastrum


									-25	-	വവ	10
[Bay of Quinte (B)]					H	64	7 0				H 6	56
мајкег		0.3			7	0.2				ω	c	0
Solitaire		Ü			9	7	0.5			Н		H
Red Chalk (main)		0.5	0.3		31	4	c	O. U	7	0.3	0.3	7 2
. Ked Chalk (east)		0.3	Н		7	2	0.5	Н	m	ഹര	0.0	0.2
Little Clear					7		0.2			0.2	0.3	
Jerry		4	Ŋ		2	7	-			0° 2° 14	4	H
Harp		0.5	4		11	-100		0.2	n		4 ⊢	2 4 4 8 8 4 8 8 9 4 8 9 9 9 9 9 9 9 9 9 9
redtsel[[u2	(mm <sub>3</sub> /m <sub>3</sub> )		4		9	17		0.3		2	0 8 8	rt rt
นอเอ	(mm				7	က	-			H	H	0 0
Dickie	max.	1			m	13	0.2	0.2		0.2	0.2	0.0
nosson		П			7	0.5				0.2	0.2	0
срир		0.5			29	ი ი	° °		-1	19	11	0 10 10 6
виск		0.3	0.2		7	1	0.2	•	<b>→</b>	0.2	0.2	0 .
Blue Chalk		1	4		Ŋ	П	0.3			0.2 13 28	0.3	0.3
bniwgia					П	-	0.2			0.3	П	↔
Basshaunt					7	2				H	0.3	0.2
[netwater]												
												9 8
	ont'd		ria				ഗ		S	Ş		phyce
	AE (C.	onas	rnacotus Planctonema Planktosphaeria	a nas	ula	Smus	Schroederia Schroederia	eldia trum	Sorastrum Sphaerocystis	Sphaerozosma Spermatozopsis Spondylosium Staurastrum	Staurodesmus Stichococcus Tetraëdron Tetrastrum Treubaria	Trochiscia Ulothrix Westella Xanthidium Unid. Chlorophyceae
	PHYCE	Pedinomonas	rnacotus Planctonema Planktospha	Polytoma Pteromonas	Quadrigula Badiococcus	Scenedesmus	Schroederia	Scourfieldia Selenastrum	Sorastrum Sphaerocy:	Sphaerozosme Spermatozope Spondylosiur Staurastrum	Staurodesmi Stichococci Tetraëdron Tetrastrum Treubaria	Trochiscia Ulothrix Westella Xanthidium Unid. Chlon
1979	CHLOROPHYCEAE (Cont'd)	Pe	26	Po Ptt	On	SC	SS	Se	Sp	Strans	T T E	T X X A L L L L L L L L L L L L L L L L L
19	1 5											


									-2	6-							
[Bay of Quinte (B)			14 991		557 54	277	211		15653	7	12		35	949	355		
Mg] ker			12		10						0.3		13		3 24		
Solitaire			11		48						0				1 74		7
Red Chalk (main)			45		385	c	2.	•	-	00.	0,3		7	0.2	39		
Red Chalk (east)			25		295	c	0		12	m,⊢l			26		64		
Little Clear			15		528 2					_	-		-		ω		
Jerry			146		64 5					· 러	4		123	٠.0	16 224		
Длен			48	,	223		19	0.2	38		7		276		58		m
reatheal[[u2	m <sub>3</sub> )	н	4		48		က	Ū	73		7		15		9 7 8		. 7
ejeu	(mm <sub>3</sub> /m <sub>3</sub> )				652	·	4				7		რ		4.0		
Dickie	max. (	,	91		273 6	, c	2 5		274	Ŋ	ന		20		6 1033 4		
Crosson	ma		29		00		0 7.		7								
Сһиь	-		73		12		) ท		23	-	9		11	Н	1 86 110 1		e 0 • 3
Bnck			89 7		ω	m	0.0		10 2				22 ]	.*	288 0.3		0
Blue Chalk			22 8		1 4	٦,	o =i				-			Н	5 110 28 0,		н
bniwei8 		е	5		13 481	,	<b>⊣</b> ო		6 0.3	H			7		3 1.		
Basshaunt		0	13 2								Н				28 1:		
			,		185												
[Tearwater]																	
					. *												
																	opsis
	CEAE	· v	D D		scus		æ	ď				a aenia	n i a ma	obia	atoms		Isthmochloron Ophiocytium Pseudopolyedriopsis
	IOPHY	Achnanthes	Asterionella Atthone	Cocconeis	Coscinodiscus Cyclotella	oma oma	Eunotia Fragilaria	rrustulla Gomphonema	sira	cula	Nitzshia	Rhoicosphaenia Rioicosphaenia	Khizosolenia Skeletonema	Stenopterobia Stephanodiscus	Surirella Synedra Tabellaria Unid. Diatoms	YCEAE	Isthmochloron Ophiocytium Pseudopolyedr
1979	BACILLARIOPHYCEAE	Achnant	Asterio	Cocc	Coscinod Cyclotel	Diatoma	Frag	Gomp	Melosira	Mavicula Navicula	Nitzshia	Rhoj	Skel	Step	Synedra Synedra Tabella Unid. D	XANTHOPHYCEAE	Isth Ophi Pseu
19	BAC															XAN	


The next 16 pages (pp.28-43) contain plots (note the logarithmic scale) of the dominant classes and genera of all 16 lakes for the May-November periods of 1976-1979. Total phytoplankton biomass is also shown and blue-green algae have been divided into two groups: bloom forming genera (Microcystis, Aphanizomenon, Coelosphaerium, Gomphosphaeria, Anabaena, Oscillatoria, Lyngbya, and Gloeotrichia) and the non-bloom forming genera.


<u>Lake</u>	Page
Blue Chalk Lake	28
Red Chalk Lake, Main Basin	29
Red Chalk Lake, East Basin	30
Harp Lake	31
Jerry Lake	32
Dickie Lake	33
Chub Lake	34
Glen Lake	35
Basshaunt Lake	36
Buck Lake	37
Solitaire Lake	38
Little Clear Lake	39
Gullfeather Lake	40
Walker Lake	41
Bigwind Lake	42
Crosson Lake	43












































## REFERENCES

- NICHOLLS, K.H. and E.C. CARNEY. 1979. The taxonomy of Bay of Quinte phytoplankton and the relative importance of common and rare taxa. Can. J. Bot. 57:1591-1608.
- NICOLLS, A., R. REID and R. GIRARD. 1983. Morphometry of the Muskoka-Haliburton Study lakes. Ont. Min. Envir. Data Report DR 83/3.
- SCHEIDER, W.A., R.A. REID, B. LOCKE and L.D. SCOTT. 1983. Studies of lakes and watersheds in Muskoka-Haliburton, Ontario:

  Methodology. Ont. Min. Envir. Data Report DR 83/1.

Ontario. Ministry of the Environment Phytoplankton of lakes in the Muskoka-Haliburton area.

1983

LABORATORY & RESEARCH LIBRARY MINISTRY OF THE ENVIRONMENT

